

TEST FOR RIGHT-HANDED b QUARK DECAYS¹

JAMES F. AMUNDSON, JONATHAN L. ROSNER, AND MIHIR P. WORAH
Enrico Fermi Institute and Department of Physics, University of Chicago
5640 S. Ellis Ave., Chicago, IL 60637, USA

and

MARK B. WISE
Lauritsen Laboratory of Physics, California Institute of Technology
Pasadena, CA 91125

ABSTRACT

Gronau and Wakaizumi have proposed a model in which the dominant b decays are due to exchange of a new right-handed gauge boson. A test of this model via the study of polarized Λ_b baryons produced in $e^+e^- \rightarrow Z \rightarrow \Lambda_b + X$ is suggested.

It is conventionally assumed that the b quark decays left-handedly. However, it turns out to be surprisingly hard to exclude the possibility [1] that the dominant b decays to charm occur via a *right*-handed coupling, as long as the coupling to leptons in such decays is also right-handed. One cannot just look at the beta-decay spectrum, which is the same for $(V - A) \times (V - A)$ and $(V + A) \times (V + A)$ couplings. Instead, one needs to study decays of polarized b quarks.

In the present report, based on the work of Ref. [2], we suggest that the reaction $e^+e^- \rightarrow b\bar{b} \rightarrow \Lambda_b + \dots$ is likely to give polarized b 's, whose decay to leptons can distinguish among models. We urge, in particular, that the LEP experiments (some of which [3, 4] have already presented evidence for Λ_b) analyze their data with the possibility of $(V + A) \times (V + A)$ couplings in mind. Some aspects of the polarization studies suggested here have been mentioned previously [5, 6].

The reaction $e^+e^- \rightarrow Z \rightarrow b\bar{b}$ is expected to give rise to b quarks with polarization $\mathcal{P} \simeq -0.93$ for a weak mixing angle $\sin^2 \theta = 0.233$. The Λ_b produced in the subsequent fragmentation of these b quarks should retain this polarization in the heavy-quark limit [6], in the absence of hard-gluon emission. The spin of the Λ_b is carried entirely by the b quark; the light quarks in it are coupled up to spin zero. If, on the other hand, the b quark ends up in a (spinless) B meson, information on its polarization is lost.

The inclusive $\Lambda_b \rightarrow \text{charm semileptonic decay}$ can be treated in the heavy-quark limit as a free-quark decay $b \rightarrow ce^- \bar{\nu}_e$. If the b has polarization $\mathcal{P}$, the expression

¹ Presented by Jonathan L. Rosner at DPF 92 Meeting, Fermilab, November, 1992.

for the normalized decay distribution in the b rest frame may be written

$$\begin{aligned} \frac{1}{\Gamma} \frac{d^2\Gamma}{dx d(\cos\psi)} &= \frac{3x^2(1-\zeta)^2}{f(m_c^2/m_b^2)} \left[1 - \frac{2}{3}x + \frac{2x-1}{3}\xi\mathcal{P}\cos\psi \right. \\ &\quad \left. + \zeta \left\{ 1 - \frac{1}{3}x + \frac{1+x}{3}\xi\mathcal{P}\cos\psi \right\} \right] , \end{aligned} \quad (1)$$

where $x \equiv 2E_e^*/m_b$, E_e^* is the electron energy in the b rest frame, ψ is the angle between the electron momentum and the spin quantization axis in this frame, $\xi = \pm 1$ for $V \pm A$ couplings at both vertices, and

$$\zeta \equiv \frac{m_c^2}{m_b^2(1-x)} \quad ; \quad f(y) \equiv 1 - 8y + 8y^3 - y^4 - 12y^2 \log y \quad . \quad (2)$$

The electron spectrum (1) is considerably harder for $\xi\mathcal{P}\cos\psi \simeq 1$ than for $\xi\mathcal{P}\cos\psi \simeq -1$, as one can also see using familiar helicity arguments.

In the reaction $e^+e^- \rightarrow Z \rightarrow \Lambda_b + X$, the b quark has a large momentum along the axis of a fairly well-defined jet. In order to obtain the signal for Λ_b production, one can select events with a lepton at some minimum transverse momentum p_T with respect to this axis and with an inclusively produced Λ baryon of the appropriate sign [3,4]. In this frame, let us imagine the b spin to be quantized along its direction of motion. The electron transverse momentum and total energy are then $p_T = (m_b x \sin\psi)/2$ and $E_e = m_b x \gamma_b (1 + v_b \cos\psi)/2$, where v_b is the b quark's velocity and $\gamma_b \equiv (1 - v_b^2)^{-1/2}$. Using these variables we may transform the distribution (1) to obtain

$$\frac{1}{\Gamma} \frac{d\Gamma}{dE_e} \Big|_{p_T \geq p_T^{\min}} = \int dx \theta(p_T - p_T^{\min}) \frac{2}{m_b x \gamma_b v_b} \frac{1}{\Gamma} \frac{d^2\Gamma}{dx d(\cos\psi)} \quad (3)$$

Examples of this distribution are shown in Fig. 1 for various values of $p_T^{\min}$ and for the two limiting cases $\xi\mathcal{P} = 1$ (close to the standard model) and $\xi\mathcal{P} = -1$ (close to the model of Ref. [1]). We have taken the b quark to have a laboratory energy of 45 GeV, so that $\gamma_b = 9$. We have taken $m_c = 1.66$ GeV, $m_b = 5$ GeV, and have ignored the electron mass. A distinction on the basis of the electron energy spectrum is clearly possible, for example, when $p_T^{\min} = 0.8$ GeV.

The distributions (1) – (3) can be convoluted with more realistic functions for fragmentation of a b quark if desired. These functions are already in use in the Monte Carlo simulations employed in Refs. [3] and [4], so it should be a simple matter to reproduce distributions analogous to those in Fig. 1 which are more appropriate to the actual experiments. The fragmentation of a b quark to a Λ_b is not actually known; the corresponding function for $b \rightarrow B$ is peaked around $p_B/p_b \approx 0.7$. A potential source of depolarization is production of Λ_b via the decay of Σ_b^* and Σ_b . If these two resonances are more closely spaced than their natural widths, they can act coherently to preserve the b quark polarization, but if they are too widely spaced, some depolarization will result. (This question has been considered in Ref. [7].) The study of the corresponding spacing between Σ_c and Σ_c^* should shed some light

on this question.

We thank M. Gronau and J. Kroll for helpful comments. This work was supported in part by the United States Department of Energy under grant No. DE AC02 90ER40560.

Figure 1: Distributions in electron laboratory energy for semileptonic decays of a b quark with laboratory energy 45 GeV and values of electron $p_T^{\text{min}} = 0, 0.8, \text{ and } 1.6$ GeV (curves in descending order). Distributions are normalized to unit area for $p_T^{\text{min}} = 0$. Solid curves: $\xi\mathcal{P} = +1$; dashed curves: $\xi\mathcal{P} = -1$.

1. M. Gronau and S. Wakaizumi, *Phys. Rev. Lett.* **68** (1992) 1992; M. Gronau, *Phys. Lett. B* **288** (1992) 90.
2. J. F. Amundson, J. L. Rosner, M. Worah, and M. B. Wise, *Phys. Rev. D* **47** (1993), to be published.
3. D. Decamp *et al.* (ALEPH Collaboration), *Phys. Lett. B* **278** (1992) 209.
4. P. D. Acton *et al.* (OPAL Collaboration), *Phys. Lett. B* **281** (1992) 394.
5. J. G. Körner and M. Krämer, *Phys. Lett. B* **275** (1992) 495.
6. T. Mannel and G. A. Schuler, *Phys. Lett. B* **279** (1992) 194.
7. F. Close, J. Körner, R. Phillips, and D. Summers, *J. Phys. G* **18** (1992) 1716.